

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 753 065 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
**14.05.2003 Bulletin 2003/20**

(51) Int Cl.7: **C12N 15/13, C07K 16/28,  
C07K 16/00**

(21) Application number: **95913225.9**

(86) International application number:  
**PCT/GB95/00692**

(22) Date of filing: **28.03.1995**

(87) International publication number:  
**WO 95/026403 (05.10.1995 Gazette 1995/42)**

### (54) **ANTIBODIES AGAINST E-SELECTIN**

ANTIKÖRPER GEGEN E-SELEKTIN

ANTICORPS CONTRE LA SELECTINE E

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL  
PT SE**

(30) Priority: **29.03.1994 GB 9406243  
29.07.1994 GB 9415331**

(43) Date of publication of application:  
**15.01.1997 Bulletin 1997/03**

(73) Proprietor: **CELLTECH THERAPEUTICS LIMITED  
Slough, Berkshire SL1 4EN (GB)**

(72) Inventors:  
• **OWENS, Raymond John  
Berkshire SL1 4EN (GB)**  
• **ROBINSON, Martyn Kim  
Berkshire SL1 4EN (GB)**

(74) Representative: **Mercer, Christopher Paul  
Carpmaels & Ransford  
43, Bloomsbury Square  
London WC1A 2RA (GB)**

(56) References cited:  
**EP-A- 0 323 806 EP-A- 0 438 312  
WO-A-88/07089 WO-A-91/09967  
WO-A-93/22436**

- **J. IMMUNOL., vol.147, no.8, 15 October 1991  
pages 2657 - 2662 LUND ET AL. 'Human  
FcgammaRI and FcgammaRII interact with  
distinct but overlapping sites on human IgG'  
cited in the application**
- **TRANSPLANT PROC., vol.22, no.4, 1990 pages  
1991 - 1993 LEUWENBERG ET AL.  
'Characterization of two monoclonal antibodies  
directed against an adhesion molecule on  
human endothelial cells' cited in the application**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 753 065 B1**

## Description

**FIELD OF THE INVENTION**

**[0001]** This invention relates to antibodies having specificity for E-selectin characterised in that said antibodies are whole antibodies of neutral isotype, to processes for preparing said antibodies, to pharmaceutical compositions containing said antibodies, and to medical uses of said antibodies.

**BACKGROUND OF THE INVENTION**

**[0002]** Selectins are a family of structurally related transmembrane glycoproteins implicated in the adhesion of leukocytes to vascular endothelial cells. The three known members, designated E-, P- and L- selectin are composed of three types of domain, an amino terminal C-type lectin domain, one EGF-like domain and between two and nine complementary regulatory repeats. Stimulation of endothelium by inflammatory cytokines e.g. IL-1, or TNF results in the upregulation of E-selectin expression on the cell surface.

**[0003]** Experiments *in vitro* have shown that E-selectin can support the adhesion of polymorphonuclear cells, monocytes and a subpopulation of T-lymphocytes (see for example, Bevilacqua *et al* (1989) Science **243** 1160-1165; Picker *et al* (1991) Nature **349** 796-799 and Leeuwenberg *et al* (1992) Scand. J. Immunol **35** 335-341). Mouse antibodies to E-selectin that block PMN binding *in vitro* have been shown to reduce extravasation of PMNs (neutrophils) in animal models (Mulligan, M. *et al*, J. Clinical Investigation **88**, 1396-1406 (1991) and Gundel, R. *et al*, J. Clinical Investigation **88**, 1407-1411 (1991)).

**[0004]** E-selectin thus appears to play a key role in the movement of leukocytes to sites of inflammation due to injury or infection. A corollary of this is that the expression of E-selectin is increased in certain inflammatory diseases. Hence E-selectin contributes to the disease process by supporting the adhesion of leukocytes which in turn cause tissue damage. It follows that an antibody to E-selectin that blocks this process would attenuate the extent or severity of the inflammation and hence be of therapeutic benefit.

**[0005]** Since most available monoclonal antibodies are of rodent origin, they are naturally antigenic in humans and thus can give rise to an undesirable immune response termed the HAMA (Human Anti-Mouse Antibody) response. Therefore, the use of rodent monoclonal antibodies as therapeutic agents in humans is inherently limited by the fact that the human subject will mount an immunological response to the antibody and will either remove it entirely or at least reduce its effectiveness.

**[0006]** Proposals have been made for making non-human MAbs less antigenic in humans using engineering techniques. These techniques generally involve the use of recombinant DNA technology to manipulate DNA sequences encoding the polypeptide chains of the antibody molecule. A simple form of engineering antibodies involves the replacement of the constant regions of the murine antibody with those from a human antibody (Morrison *et al* (1984) Proc. Natl. Acad. Sci. USA **81** 6851-55; Whittle *et al* (1987) Prot. Eng. **1** 499-505). The lowering of the level of the HAMA response to the chimeric antibodies leads to the expectation that further engineering of the variable region outside of the antigen binding site may abolish the response to these regions and further reduce any adverse response.

**[0007]** A more complex form of engineering of an antibody involves the redesign of the variable region domain so that the amino acids constituting the murine antibody binding site are integrated into the framework of a human antibody variable region. This has led to the reconstitution of full antigen binding activity in a number of cases (Co *et al* (1990) J. Immunol. **148** 1149-1154; Co *et al* (1992) Proc. Natl. Acad. Sci. USA **88** 2869-2873; Carter *et al* (1992) Proc. Natl. Acad. Sci. **89** 4285-4289; Routledge *et al* (1991) Eur. J. Immunol. **21** 2717-2725 and International Patent Specifications Nos. WO 91/09967; WO 91/09968 and WO 92/11383).

**[0008]** Naturally occurring and engineered human antibodies may be regarded as bifunctional agents, with the N-terminal variable region responsible for antigen binding and sequences within the C-terminal part responsible for determining interactions with the various cell types which participate in immune responses. Recognition of these effector sites on antibodies by specific cell surface receptors on cytotoxic cells can result in antibody-dependent cellular cytotoxicity and complement mediated lysis. This can result in killing of the cell presenting the antigen.

**[0009]** E-selectin is expressed on the surface of endothelial cells. The loss of endothelial cells as a result of antibody bound to target antigen is highly undesirable. Endothelial cells make up the endothelium which forms a barrier between the tissues of the body and the vascular system. The loss of or damage to the structural integrity of the endothelium is extremely disadvantageous and can lead to oedema and vasculitis. It is highly advantageous therefore to avoid depletion of the endothelial cell population while blocking the target antigen. Recent papers by Podolsky *et al* (J. Clin. Invest. **92** (1993) 372-380), Westphal *et al* Clin. Exp. Immunol. **96** 444-449 (1994), and the Editorial Lancet **337** (1991) confirm that the use of whole antibody is undesirable due to undesirable effector functions mediated via the Fc region of the antibody. Another group has attempted to overcome the problem of undesirable effector functions by the use of antibody fragments lacking the effector signals which result in antibody-dependent cellular cytotoxicity (Mulligan *et al*,

J. Clinical Investigation 88, 1396-1406 (1991)). Antibody fragments are known, however, to have a short half-life (Pimm *et al* Nuclear Medicine Communication 10, 585-593 (1989); Molthoff *et al* Br. J. Cancer 65, 677-683 (1992) and Buist *et al* Cancer Res. 53 5413-5418 (1993)) making their therapeutic usefulness in the treatment of many diseases extremely limited.

## DESCRIPTION OF ASPECTS OF THE INVENTION

[0010] The present invention provides a novel solution to this problem of preventing depletion of the target endothelial cell population. We have found that by making a whole anti-E-selectin antibody of neutral isotype it is possible to produce a therapeutically useful antibody which does not result in endothelial cell depletion. In a first aspect the invention therefore provides an antibody having a specificity for E-selectin wherein the complementarity determining regions of the variable domains in said antibody correspond to those of the murine antibody ENA-2 and the following framework residues in said antibody correspond to the same residues of the murine antibody ENA-2:

- heavy chain: 48, 67, 69, 73, 93 & 94
- light chain: 48, 60, 63, 70, 111 & 113

[0011] Anti-E-selectin antibody may be prepared using well-known immunological techniques employing E-selectin as antigen. Any suitable host may, for example, be immunised with E-selectin or activated HUVEC (human umbilical vein endothelial cells) and splenocytes of lymphocytes recovered and immortalised using for example the method of Kohler *et al*. Eur. J. Immunol. 6, 511 (1976). The resulting cells are diluted and cloned to obtain a single genetic line producing anti-E-selectin antibodies in accordance with conventional practice. Where it is desired to produce recombinant anti-E-selectin antibodies these may be produced using methods well known in the art.

[0012] ENA-2 is a mouse IgG1/kappa antibody that binds to the lectin/EGF region of human E-selectin and blocks cell binding. (Leeuwenberg *et al* (1990) Transplantation Proceedings 22 (4) 1991-1993).

[0013] In a second aspect of the invention, the antibody of the first aspect is of neutral isotype. The approach of using a whole antibody of neutral isotype has not been tried before in this area. The term "neutral isotype" means that the interactions with antibody Fc receptors i.e. FcRI, FcRII and FcRIII and complement are absent or so weak as to cause minimal detrimental physiological effects such as antibody dependent cellular cytotoxicity (ADCC) and/or complement mediated lysis and also the antibody produces a minimal immune response in the host. As used herein, the term "minimal immune response" is used to denote a typical primate immune response to an IV injection of a human or engineered human antibody.

[0014] Several regions of the Fc region of antibodies have been implicated in modulating effector functions (see for example European Patent No. 307434B and Lund *et al* (1991) J. Immunot. 147 2657-2662). For example Lund *et al* (1991) and other groups have implicated Leu 235 in the CH2 domain of human IgG3 heavy chain in binding of antibody to the high affinity receptor on mononuclear phagocytes (FcRI). Thus by altering this residue it is possible to produce an antibody lacking FcRI binding activity.

[0015] In a further aspect the invention provides an antibody characterised in that said antibody is a whole antibody of neutral isotype having specificity for E-selectin wherein one or more amino acid residues in the Fc region of said antibody has been altered from that in the naturally occurring sequence.

[0016] In a preferred embodiment of this aspect the invention therefore provides an antibody characterised in that said antibody is a whole antibody of neutral isotype having specificity for E-selectin wherein one or more amino acid residues in the Fc region of said antibody including residue 235 in the CH2 domain has been altered from that in the naturally occurring sequence.

[0017] Residue 235 occurs in the N-terminal region of the CH2 domain of the heavy chain. We have found altering the naturally occurring residue e.g. residue Leu 235 to an alanine is particularly advantageous since the conservative nature of the amino acid change is less likely to produce undesirable structural changes in the molecule, which may result in the antibody being immunogenic.

[0018] In a preferred embodiment the antibody of the invention has an alanine residue at position 235 in the CH2 domain and in a particularly preferred embodiment the antibody has a human  $\gamma 4$  isotype.

[0019] Alteration of antibody side chain interaction with FcR1 receptor may similarly be achieved by replacing leucine with isoleucine, valine, threonine or glutamic acid. It will be readily apparent to one skilled in the art that a number of other substitutions are possible within the overall aim of not introducing any undesirable structural changes in the molecule, which may lead to immunogenicity.

[0020] Where the antibody has isotypes such as human  $\gamma 1$ ,  $\gamma 2$  or  $\gamma 3$  it will be apparent to one skilled in the art that further alterations to amino acid residues will be required to alter FcRI, FcRII and FcRIII binding where appropriate and also to minimise complement fixation. For example, alteration of residues 235 and 234 of the heavy chain of  $\gamma 1$ , and  $\gamma 3$  antibodies is known to affect FcRI and FcRII binding (Burton and Woof *Adv. Immunol* (1992) 1: Academic

Press), and similarly amino acid residues 234, 235, 330, 331, 318, 320 and 322 have been shown to be involved in binding and activation of complement (Xu *et al* (1994) J. Biol. Chem. 269 (5) 3469-3474, Published European Patent No. EP 307434B and published International Patent Application No. WO 9429351). See also, Woof *et al* Mol. Immunol 23 319-330 (1986), Burton *et al* Nature 288 338-44 (1980); Burton Mol. Immunol 22 161-206 (1988); Leatherbarrow *et al* Mol. Immunol 22 407-415 (1985); and Duncan *et al* Nature 332 563-4 (1988). The antibodies of the invention preferably have a human isotype.

[0021] The standard techniques of molecular biology may be used to prepare DNA sequences coding for the antibodies according to the invention. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate.

[0022] Suitable processes which may be used to alter the residue at position 235 include the PCR strand overlap procedure PCR mutagenesis, as described for example in the teaching of PCR Technology Principles and Applications for DNA Amplification (1989), Ed. H.A. Erlich, Stockton Press, N.Y., London, and oligonucleotide directed mutagenesis (Kramer *et al*, (Nucleic. Acid. Res. 12, 9441 (1984)). Suitable methods are also disclosed in Published European Patent No. EP307434B.

[0023] The alteration at position 235 or any other position of the molecule may be introduced at any convenient stage in the antibody production process. For example, where the antibody is a CDR-grafted antibody, the change may be made before, or more conveniently after CDR-grafting has been completed. This is described in more detail in the accompanying examples.

[0024] In a preferred embodiment the antibody molecule of the invention is an IgG and most preferably has a human  $\gamma 4$  isotype.

[0025] It has further been found (Angal *et al* (1993) Molecular Immunol 30 105-108) that the sequence of the hinge region of antibodies of the  $\gamma 4$  isotype, i.e. Cys-Pro-Ser-Cys can give rise to alternative forms of the antibody in association with correctly folded and assembled forms. This can be overcome by altering the Ser residue at position 228 to a Pro residue using for example site directed or oligonucleotide directed mutagenesis. The anti-E selectin antibodies of the invention which are of  $\gamma 4$  isotype preferably have the sequence Cys-Pro-Pro-Cys at the hinge region, and most preferably also have an alanine residue at position 235 in the CH2 domain.

[0026] The residue numbering used herein is according to the EU index described in Kabat *et al* [(1991) in: Sequences of Proteins of Immunological Interest, 5th Edition, United States Department of Health and Human Services].

[0027] The term engineered human antibody molecule is used to describe a molecule having an antigen binding site derived from an immunoglobulin from a non-human species, the remaining immunoglobulin-derived parts of the molecule being derived from a human immunoglobulin. The antigen binding site may comprise either complete variable regions fused onto human constant domains or only the complementarity determining regions (CDRs) grafted onto appropriate human framework regions in the variable domains.

[0028] The human immunoglobulin derived parts of the engineered human antibody molecule may be derived from any suitable human immunoglobulin. For instance where the engineered human antibody molecule is a CDR-grafted antibody molecule, appropriate variable region framework sequences may be used having regard to class/type of the donor antibody from which the antigen binding regions are derived. Preferably the type of human framework used is of the same/similar class/type as the donor antibody. Advantageously the framework is chosen to maximise/optimize homology with the donor antibody sequence particularly at positions spacially close or adjacent to the CDRs. Examples of human frameworks which may be used to construct CDR-grafted antibodies are LAY, POM, TUR, TEI, KOL, NEWM, REI and EU; for instance KOL and NEWM for the heavy chain and REI for the light chain and EU for both the heavy chain and light chain.

[0029] In a preferred method the human frameworks are chosen by comparing the sequences of the donor and acceptor heavy and light chains and choosing the human framework sequence which is most homologous to the donor antibody.

[0030] The ENA-2 Vh domain shows closest sequence homology to group 1 human heavy chains and consequently the group 1 human antibody Eu was chosen as the framework for both the heavy and light chain variable domains.

[0031] The light or heavy chain variable domains of the engineered human antibody molecule may be fused to human light or heavy chain constant domains as appropriate, (the term Fc region and 'heavy chain constant domains' as used herein are to be understood to include hinge regions unless specified otherwise). The human constant domains of the engineered human antibody molecule, where present, may be selected having regard to the proposed function of the antibody, in particular the lack of effector functions which may be required.

[0032] For example, the heavy chain constant domains fused to the heavy chain variable region may be human IgA, IgG or IgM domains. Preferably human IgG domains are used. Depending on the choice of human constant domains it may be necessary to alter specific amino acid residues to remove any undesirable effector function in order to produce an antibody of neutral isotype by, for example, using site directed or oligonucleotide directed mutagenesis. Light chain human constant domains which may be fused to the light chain variable region include human Lambda or human

Kappa chains.

**[0033]** Analogues of human constant domains may alternatively be advantageously used. These include those constant domains containing one or more additional amino acids than the corresponding human domain or those constant domains wherein one or more existing amino acids of the corresponding human domain has been substituted, added, deleted or altered. Such domains may be obtained, for example, by oligonucleotide directed mutagenesis.

**[0034]** Also human constant region domains of the engineered human antibody molecule may be selected having regard to the neutral isotype required for the antibody as defined previously.

**[0035]** By appropriate choice of immunoglobulin isotype it is possible to produce an antibody where antibody dependent complement fixation and where interaction with FcRI, FcRII and FcRIII are minimised e.g. by choosing a human  $\gamma 4$  isotype.

**[0036]** The invention further provides a process for the production of an antibody having specificity for E-selectin characterised in that said antibody is a whole antibody of neutral isotype which process comprises:

a) producing in an expression vector an operon having a DNA sequence which encodes said antibody heavy or light chain;

b) producing in an expression vector an operon having a DNA sequence which encodes a complementary antibody heavy or light chain;

c) transfecting a host cell with both operons and

d) culturing the transfected cell line to produce the antibody.

**[0037]** According to a preferred embodiment of this aspect of the invention there is provided a process for producing an engineered human antibody having specificity for E-selectin characterised in that said antibody is a whole antibody of neutral isotype which process comprises:

a) producing in an expression vector an operon having a DNA sequence which encodes an antibody heavy or light chain comprising a variable domain wherein at least one of the CDRs of the variable domain corresponds to a non-human immunoglobulin and the remaining immunoglobulin-derived parts of the antibody chain are derived from a human immunoglobulin;

b) producing in an expression vector an operon having a DNA sequence which encodes a complementary antibody light or heavy chain comprising a variable domain wherein at least one of the CDRs of the variable domain corresponds to a non-human immunoglobulin and the remaining immunoglobulin-derived parts of the antibody chain are derived from a human immunoglobulin;

c) transfecting a host cell with both operons; and

d) culturing the transfected cell line to produce the engineered human antibody molecule.

**[0038]** The CDRs of the variable domain correspond to the same non-human immunoglobulin which is ENA-2.

**[0039]** In a particularly preferred embodiment of this aspect of the invention, at least one of the expression vectors contains a DNA sequence encoding an antibody heavy chain wherein one or more amino acid residues in the Fc region of said antibody most preferably including residue 235 in the CH2 domain has been altered from that in the naturally occurring sequence

**[0040]** The change at amino acid residue 235 or at any other position may also be made after the whole antibody has been assembled using techniques such as site directed mutagenesis.

**[0041]** The cell line may be transfected with two vectors, the first vector containing the operon encoding the light chain-derived polypeptide and the second vector containing the operon encoding the heavy chain derived polypeptide. Preferably the vectors are identical except in so far as the coding sequences and selectable markers are concerned so as to ensure as far as possible that each polypeptide chain is equally expressed.

**[0042]** Alternatively, a single vector may be used, the vector including a selectable marker and the operons encoding both light chain- and heavy chain-derived polypeptides.

**[0043]** In further aspects, the invention also includes DNA sequences coding for the heavy and light chains of the antibodies of the present invention, cloning and expression vectors containing these DNA sequences, host cells transformed with these DNA sequences and processes for producing the heavy or light chains and antibody molecules comprising expressing these DNA sequences in a transformed host cell.

[0044] The general methods by which the vectors may be constructed, transfection methods and culture methods are well known *per se* (see for example Maniatis *et al* (1982) (Molecular Cloning, Cold Spring Harbor, New York) and Primrose and Old (1980) (Principles of Gene Manipulation, Blackwell, Oxford) and the examples hereinafter).

[0045] The DNA sequences which encode the ENA-2 heavy and light chain variable domain amino acid sequences (and the corresponding deduced amino acid sequences) are given hereafter in Figure 1.

[0046] DNA coding for human immunoglobulin sequences may be obtained in any appropriate way. For example, amino acid sequences of preferred human acceptor frameworks such as, LAY, POM, KOL, REI, EU, TUR, TEI and NEWM are widely available to workers in the art.

[0047] The standard techniques of molecular biology may be used to prepare DNA sequences coding for CDR-grafted products. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate. For example, oligonucleotide directed synthesis (Jones *et al* (1986) Nature 321 522-525). Also oligonucleotide directed mutagenesis of a pre-existing variable domain region (Verhoeyen *et al* (1988) Science 239 1534-1536; Reichmann *et al* (1988) Nature 332 323-327).

[0048] Enzymatic filling-in of gapped oligonucleotides using T4 DNA polymerase (Queen *et al* (1989) Proc. Natl. Acad. Sci. USA 86 10029-10033; Published International Patent Application No. WO 90/07861) may be used.

[0049] Any suitable host cell/vector system may be used for expression of the DNA sequences coding the antibody heavy and light chains e.g. for the chimeric or CDR-grafted heavy and light chains. Bacterial e.g. E.coli and other microbial systems may be used. Eucaryotic e.g. mammalian host cell expression systems may also be used to obtain antibodies according to the invention, particularly for production of larger chimeric or CDR-grafted antibody products. Suitable mammalian host cells include CHO cells and myeloma or hybridoma cell lines, for example NSO cells. NSO cells are particularly preferred.

[0050] In the engineered human antibody according to the invention, the heavy and light chain variable domains may comprise either the entire variable domains of a non-human antibody such as the murine antibody ENA-2, or may comprise framework regions of a human variable domain having grafted thereon one, some or all of the CDRs of a non-human antibody such as the murine antibody ENA-2. Thus the engineered human antibody may comprise a chimeric engineered human antibody or a CDR-grafted engineered human antibody.

[0051] When the engineered human antibody is a CDR-grafted antibody, in addition to the CDRs, specific variable region framework residues may be altered to correspond to non-human e.g. ENA-2 mouse residues. Preferably the CDR-grafted antibodies of the present invention include CDR-grafted antibodies as defined in our Published International Patent Specification No. WO91/09967.

[0052] Preferably the CDRs of the heavy chain correspond to the Kabat defined MAb ENA-2 CDRs at all of CDR1 (31 to 35), CDR2 (50 to 65) and CDR3 (95 to 102). In addition the heavy chain may have mouse ENA-2 residues at one or more of residues 48, 67, 69, 73, 93 and 94. Similarly the light chain may have mouse ENA2 residues at positions 48, 60, 63, 70, 111 and 113.

[0053] The present invention also provides therapeutic and diagnostic compositions comprising the antibodies of the invention and the uses of these products and the compositions in therapy and diagnosis. Such compositions typically comprise an antibody according to the invention together with a pharmaceutically acceptable excipient, diluent or carrier, e.g. for *in vivo* use.

[0054] Thus in a further aspect the invention provides a therapeutic or diagnostic composition comprising an antibody according to the invention in combination with a pharmaceutically acceptable excipient, diluent or carrier.

[0055] The invention also provides a process for the preparation of a therapeutic or diagnostic composition comprising admixing an antibody according to the invention together with a pharmaceutically acceptable excipient, diluent or carrier.

[0056] The antibody may be the sole active ingredient in the therapeutic or diagnostic composition or may be accompanied by one or more other active ingredients. The therapeutic and diagnostic compositions may be in unit dosage form, in which case each unit dose comprises an effective amount of the antibody of the invention.

[0057] Furthermore, the invention also provides methods of therapy and diagnosis comprising administering an effective amount of an antibody according to the invention to a human or animal subject.

[0058] The antibodies and compositions may be for administration in any appropriate form and amount according to the therapy in which they are employed.

[0059] The therapeutic or diagnostic composition may take any suitable form for administration, and, preferably is in a form suitable for parenteral administration e.g. by injection or infusion, for example by bolus injection or continuous infusion. It may for example be administered intravenously, intramuscularly, intradermally or intraperitoneally. Where the product is for injection or infusion, it may take the form of a suspension, solution or emulsion in an oily or aqueous vehicle and it may contain formulatory agents such as suspending, preservative, stabilising and/or dispersing agents.

[0060] Alternatively, the antibody or composition may be in dry form, for reconstitution before use with an appropriate sterile liquid. The antibody may also be formulated for topical administration.

[0061] If the antibody or composition is suitable for oral administration, the formulation may contain, in addition to the active ingredient, additives such as: starch - e.g. potato, maize or wheat starch or cellulose - or starch derivatives such as microcrystalline cellulose; silica; various sugars such as lactose; magnesium carbonate and/or calcium phosphate. It is desirable that, if the formulation is for oral administration it will be well tolerated by the patient's digestive system. To this end, it may be desirable to include in the formulation mucus formers and resins. It may also be desirable to improve tolerance by formulating the antibody or compositions in a capsule which is insoluble in the gastric juices. It may also be preferable to include the antibody or composition in a controlled release formulation.

[0062] The antibodies of the invention may be used for the treatment of a human or animal subject suffering from or at risk of a disorder associated with increased E-selectin expression the method comprising administering to the subject an effective amount of the antibody or composition of the invention. In particular, the human or animal subject may be suffering from an inflammatory disorder such as a skin disorder e.g. psoriasis.

[0063] The antibodies of the invention are particularly useful in the treatment of inflammatory diseases generally. They are particularly useful in the treatment of inflammatory skin diseases e.g. psoriasis, contact dermatitis and eczema; inflammatory bowel disease e.g. Crohn's disease and ulcerative colitis, in lung inflammatory disorders, e.g. ARDS; arthritis, e.g. rheumatoid arthritis; vasculitis, liver disease e.g. alcoholic hepatitis and cirrhosis; and thermal trauma.

[0064] Therapeutic and diagnostic uses typically comprise administering an effective amount of an antibody according to the invention to a human subject. The exact dose to be administered will vary according to the use of the antibody and on the age, sex and condition of the patient but may typically be varied from about 0.1mg to 1000mg for example from about 1mg to 500mg. The antibody may be administered as a single dose or in a continuous manner over a period of time. Varying doses may be repeated as appropriate. The antibody may be formulated in accordance with conventional practice for administration by any suitable route and may generally be in a liquid form (e.g. a solution of the antibody in a sterile physiologically acceptable buffer) for administration by for example an intravenous, intraperitoneal or intramuscular route.

[0065] Since the whole antibodies are of neutral isotype the interaction with Fc receptors will be minimal. This has the effect that such an antibody should block greater than 80% of human neutrophil binding to E-selectin in an *in vitro* assay and furthermore that this may be observed irrespective of the FcR status of the donor. We believe this may be a considerable advantage in that the antibody is suitable for administration to all patients thereby avoiding the necessity of determining the FcR status of the patient prior to administration of the antibody.

[0066] The antibodies according to the invention in *in vitro* assays show minimal binding to FcR1 carrying cells that do not express E-selectin thereby minimising the potential for ADCC.

[0067] The antibody according to the invention may also be used for inflamed site-specific delivery of drugs, nucleic acid and proteins and fragments thereof, radionuclides or chelated metals and other therapeutic agents. The therapeutic agents may be linked directly to the antibody or via a carrier such as for example, a liposome, virus or viral particle where the therapeutic agent to be delivered is incorporated as part of the carrier. This technology is well known in the art. Similarly the antibodies of the invention may also be used diagnostically in the identification of areas of inflammation. The antibodies may be unlabelled or may be labelled for example with a radiolabel, e.g. a radionuclide; a chelated metal; a photochemical reagent, e.g. a fluorescent compound; a dye, or a label detectable via reaction with an enzyme, substrate or cofactor, or a compound detected via NMR or ESR spectroscopy.

#### BRIEF DESCRIPTION OF THE FIGURES

[0068] The present invention is now described by way of example only, by reference to the accompanying drawings in which:

Figure 1a shows the DNA and amino acid sequence of ENA-2 VI.

Figure 1b shows the DNA and amino acid sequence of ENA-2 Vh.

Figure 2a shows the DNA and amino acid sequence of CDR-grafted ENA-2 Vh.

Figure 2b shows the DNA and amino acid sequence of CDR-grafted ENA-2 VI.

Figure 3a shows a graph of the competition binding activity of chimeric ENA-2.

— mouse ENA2  
 - - - chimeric ENA2

Figure 3b shows a graph of the competition binding activity of CDR-grafted ENA-2 antibody.

— mouse ENA2  
 - -X- - hENA-2

Figure 4a shows a graph of the cell blocking activity of chimeric and mouse ENA-2 antibodies.

— ENA2 1G  
 - - ◆ - - chimeric

Figure 4b shows a graph of the cell blocking activity of CDR-grafted and mouse ENA-2 antibodies.

—●— ENA2 FAB2

...●... ENA2 1G

-●- hENA-2

Figure 5 shows the sequence of mutagenic oligonucleotides.

Figure 6 is a histogram analysis showing the cell binding activity of CDR-grafted hENA-2 (wt) and hENA-2(L235A).

■ gamma 4

▨ no ab

▩ gamma 4 L235>A

□ gamma 4

▤ no ab

▥ gamma 4 L235>A

Figure 7 is a schematic diagram of the expression vector pENA215.

Figure 8 is a schematic diagram of the expression vector pRO102

Figure 9 is a schematic diagram of the vectors pMRR015 and pMRR011

Figure 10 shows a graph of the effect of hENA-2(L235A) on IL-1 induced PMN infiltration in baboon skin

■ saline injected sites (n=12)

▲ hENA2(L235A) treated (n=6)

▼ saline control treated (n=6)

Figure 11 shows a graph of the effect of hENA-2(L235A) on TNF induced PMN infiltration in baboon skin

■ saline injected sites (n=12)

▲ hENA2(L235A) treated (n=6)

▼ saline control treated (n=6)

P < 0.05

Figure 12 shows a graph of the effect of ENA-2 (Fab)<sub>2</sub> and controls on PMN in baboon skin after IL-1 induction.

—◇— IL-1 0.1μg (control)

—△— IL-1 0.1μg (treated)

—◇— Saline

Figure 13 shows a graph of the effect of ENA-2 (Fab)<sub>2</sub> and controls on PMN in baboon skin after TNF induction

—X— TNF 5μg (control)

—◇— TNF 5μg (treated)

—△— Saline combined

P < 0.05

Figure 14 shows a graph of circulating hENA-2(L235A) levels in the animals used in figures 12 and 13 as measured by a cell-based assay system.

Figure 15 shows a graph of white blood cell trafficking in human skin xenografts on SCID mice after intradermal challenge with:

BSA ▨

TNF ■

and i.v. treatment with hENA-2(L235A, S228P).

Figure 16 shows the effect of hENA-2(L235A)

hENA-2(L235A, S228P)

MOPC 21

on the binding of human PMN to E-selectin transfected CHO cells.

Figure 17 shows the results of an experiment to measure complement dependent lysis

1A - Total lysis using 10% SDS.

1B - Medium alone.

1C - Medium + complement.

1D - W6/32 + complement.

1E - hENA-2(L235A, S228P) + complement.

1F - hENA L235A) + complement.

2A - W6/32 + heat inactivated complement.

2B - hENA-2(L235A, S228P) + heat inactivated complement.

2C - hENA-2(L235A) + heat inactivated complement.

2D - hENA-2(L235A) + medium alone.

All error bars are ± 1 standard deviation.

Figure 18 shows micrographs of haematoxylin and eosin stain sections from biopsies of baboon skin with cytokine



induced inflammation

A) saline treated animal

B) hENA-2(L235A) treated animal.

Figure 19 shows the inhibition of binding of human neutrophils to human E-selectin transfected CHO cells

—■— ENA2 IG  
 .....●..... ENA2 FAB2  
 - - - - -●- - - - - Chimeric ENA-2

## DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

### EXAMPLE 1

#### Production of mouse ENA-2 and ENA-2F(ab')<sub>2</sub>

[0069] The hybridoma producing mouse ENA-2 was grown in DMEM medium containing 10% foetal calf serum. Antibody was purified from cell supernatants by affinity chromatography on Protein A-sepharose. This material was used to prepare F(ab')<sub>2</sub> fragments by digestion with bromelain followed by Protein A - chromatography to remove the Fc portion and undigested antibody followed by DEAE ion-exchange chromatography to remove the protease.

[0070] The first 19 amino acid residues of the ENA-2 light chain were determined by N-terminal protein sequencing.

#### Cloning mENA-2 Variable Domain Genes and Construction of Chimeric Antibodies

[0071] The heavy and light chain variable domains of ENA-2 were cloned using the polymerase chain reaction (PCR). The primers used for the heavy chain were the same as described by Jones and Bendig (1991) (Bio Technology 9 88-89) with minor modifications to enable fragments to be cloned directly into Celltech expression vectors. PCR reactions were carried out using first strand cDNA (PCR conditions 94° 1 min; 55°C 1 min; 72°C 1 min for 30 cycles) produced from total RNA by reverse transcriptase. The leader sequence primed product was inserted into pMR014 to produce the chimeric γ4 heavy chain vector pENA202. Four independent heavy chain clones were sequenced and shown to be identical except in the leader sequence as shown in Figure 1a. The ENA-2 light chain variable domain (VI) was isolated by PCR using a 5' oligonucleotide primer based on the N-terminal sequence of the murine light chain (Figure 1b) and a framework 4 3' consensus primer. The PCR product was inserted into the vector pR0102 which contains a light chain leader sequence, thus reconstructing the 5' end of VI gene. Sequencing of four independent VI clones only revealed two conservative changes in the DNA sequence (figure 1b). The reconstructed VI gene was inserted into the vector pMR15.1 to produce the chimeric Kappa light chain vector, pEN201.

#### Design and Assembly of CDR-Grafted ENA-2

[0072] The ENA-2 Vh domain showed closest sequence homology to group 1 human heavy chains. Consequently the CDR grafted version of ENA-2 was based on the group 1 human antibody Eu. The design of the CDR-grafted ENA-2 followed the criteria identified in WO 91/09967. The CDRs of both Vh and VI were defined according to Kabat *et al.* (1987) (Sequences of proteins of immunological interest 4th edition. Washington DC: United States, Department of Health and Human Services). The CDR sequences in Eu were changed to the corresponding ones of mENA-2. For the CDR grafted Vh, six framework residues were changed to the corresponding residues from mENA-2 at Met 48 lie, Val67 Ala (CDR2 packing residues contacting residue 63) lie 69 Leu, Glu 73 Lys (potential CDR2 contacting residues), Ala93 Thr (Vh/VI interface residues) and Gly 94 Val (CDR3 packing residue). In addition the following framework residues in Eu were changed to those of the human consensus sequence, Glu103Trp, Tyr104Gly, Asn105Gln and Gly107Thr. The DNA sequence of the CDR-grafted Vh gene is shown in Figure 2a. In designing the CDR grafted light chain, six changes were made to the Eu framework, Met48Ile, Ser60Asp Ile63Thr, Glu70Asp, Val111Ile Gly113 Arg. At all these positions the ENA-2 framework residue is more typical of human VI Antibody sequence than the Eu residue.

[0073] The sequence of the CDR-grafted ENA-2 VI gene is shown in Figure 2b. The CDR-grafted variable domain genes were assembled by PCR (Daugherty *et al.* (1991) Nucl. Acids Res. 19 2471-2476) using 1pmole of each assembly oligonucleotide (PCR conditions; 94°C 1 min, 55°C 1 min, 72°C 1 min for 30 cycles). CDR grafted Vh and VI genes were inserted into the vectors pMR011, pMR014 and pMR10.1 to produce the human γ1 heavy chain vector, pENA212, γ4 heavy chain vector pENA206 and the human kappa chain vector pENA204. pMR10.1 is a kappa light chain vector which lacks the Ig/terminator sequence in pMR15.1 (Figure 9).

### **Transient Expression of Chimeric and CDR-Grafted ENA-2 Antibodies**

[0074] Heavy and light chain genes were co-expressed transiently in CHO-L761 cells (Cockett *et al.*, 1991, Nucl. Acids Res. 19, 319-325) and cell supernatants assayed for assembled antibody by ELISA.

[0075] The assembly ELISA for quantifying antibody yields used microwell plates coated with a goat F(ab')<sub>2</sub> anti-human IgG Fc. Following incubation with transfected culture supernatants, bound chimeric or CDR-grafted antibody was revealed with a horse radish peroxidase (HRP)-conjugated murine anti-human kappa chain antibody using tetramethyl benzidine (TMB) as the substrate. Concentrations of chimeric or CDR-grafted whole antibody in the samples were interpolated from a calibration curve generated from serial dilutions of purified human IgG4 or IgG1 antibody standards.

### **Purification and Binding Activity of Chimeric and CDR-Grafted ENA-2 Antibodies**

[0076] Recombinant antibodies were purified from cell supernatants by affinity chromatography on Protein A-sepharose. Antigen binding activity was compared to ENA-2 in both cell blocking and competition assays. The cell blocking assay was carried out as follows:

Purified Human Polymorphs Were Pre-Incubated With Anti-CD18 Antibody and Cooled to 4°C to Minimise Integrin Binding.

Cells (4x 10<sup>5</sup> Cells/Well) were Added to CHO Cell Line Transfected with the Gene for Human E-Selectin and Incubated for 60 min at 4°C in Presence or Absence of Anti-E-Selectin Antibodies.

Cells were Washed to Remove Unbound Cells.

Adherent Cells were Lysed and Endogenous Myeloperoxidase Activity in Polymorphs was Measured in a TMB Assay.

[0077] The competition binding assay was carried out as follows.

[0078] The non-blocking anti-E-selectin 34.27 was coated onto Immunosorp 96 well plates in 0.1M NHCO<sub>3</sub> (4°C overnight at 5 µg/ml). The plate was blocked with 1% BSA (1h, room temperature) and then the Lectin/EGF fragment from E-selectin added at 50 ng/ml (100µl/well in PBS room temperature for 1h). 50µl of test sample was then added followed by 50 µl of biotinylated murine ENA-2 whole antibody. The assay was left at room temperature for 1h, streptavidin peroxidase was added and allowed to incubate at room temperature for a further 30 min. Finally, the plate was washed and colour developed with the standard TMB reagent.

[0079] In both assays chimeric and CDR-grafted ENA-2 showed similar potency to the mouse antibody. Representative data are shown in Figures 3 and 4.

[0080] The affinity of both mouse and CDR-grafted ENA-2 for E-selectin were measured using a BIAcore TH biosensor (Pharmacia). This instrument combines miniaturised fluid delivery with an optical detection system based on surface plasmon resonance, to monitor protein interactions in real time (Chaiken *et al.*, (1991) Anal. Biochem 201 197-210). The antibodies were immobilised at similar densities onto the sensor surface and the lectin/EGF fragment of E-selectin used as antigen in the fluid phase.

[0081] Assays of both antibodies were run at the same time using the same dilutions of antigen. The KD measurements are given below.

KD Measurements		
	KD (Average)	Duplicates
ENA-2	5.43 x 10 <sup>-9</sup> M	(5.38 and 5.48)
hENA2	7.98 x 10 <sup>-9</sup> M	(7.8 and 8.17)

[0082] The results indicate that the CDR-grafted ENA-2 retains approximately 70% of the affinity of the mouse antibody.

### **Construction and Evaluation of Human $\gamma$ 4 Heavy Chain Fc Mutants**

[0083] Lund *et al.*, 1991 have implicated Leu 235 in the C $\gamma$ 2 domain of human IgG3 heavy chain in binding of antibody

to the high affinity receptor on mononuclear phagocytes (FcRI). Thus changing Leu 235 to Ala or Glu drastically reduces FcRI binding activity (by approx. 90% and 100% respectively). In order to produce a version of human IgG4 lacking FcRI binding activity Leu 235 in the  $\gamma 4$  heavy chain was changed to Ala by the PCR strand overlap procedure (Ho, S. N. *et al.*, 1989), the sequences of the mutagenic oligonucleotides are given in Figure 5. Ala occurs at position 235 in human  $\gamma 2$  which does not bind to FcRI. This relatively conservative substitution was chosen to minimise the impact of the mutation on the immunogenicity of the IgG4 antibody.

[0084] A CDR grafted heavy chain vector containing the L235A mutation (pENA211) was constructed and transiently co-expressed with the CDR-grafted light chain gene in CHO L761 cells. Antibody purified from cell supernatants was evaluated for both antigen-binding and cell binding via FcRI. Competition assays showed that alteration of the Fc had no effect on the antigen-binding activity of the CDR-grafted ENA-2. The binding of hENA-2 to THP1 cells which express FcRI and JY cells which do not, and neither of which express E-selectin, were compared to hENA-2 IgG4 L235A as follows. Cells ( $5 \times 10^6$ /ml) were incubated for 1h at room temperature with either hENA-2 wt or hENA-2 L235A antibodies serially diluted in culture medium containing 10% foetal calf serum. The cells were washed with phosphate buffered saline/1% bovine serum albumin, and then incubated with a fluorescein labelled goat anti-(human IgG Fc) antibody for a further 1h at room temperature. After washing the cells to remove unbound antibody, the binding of FITC labelled antibody to the cells was detected in a FACScan analyser (Becton Dickinson). The results shown in Figure 6 confirm that the L235A mutation has removed the cell binding activity mediated by FcRI from the hENA-2 antibody.

#### Expression of CDR-grafted ENA-2 Antibodies in NSO Cells

[0085] hENA-2 (L235A) was expressed in NSO cells to produce sufficient quantities of antibody for functional characterisation in animal models. Cell lines were established using the glutamine synthetase selectable marker (See published International Patent No. WO 87/04462). A double gene expression vector was constructed by inserting the CDR grafted heavy chain gene on a Not-Bam restriction fragment into the CDR-grafted light chain vector pENA204 to produce plasmid pENA215 (Figure 7). NSO cells were transfected by electroporation and cell lines producing recombinant antibody selected using glutamine free media.

#### Construction and Expression of Human $\gamma 4$ Hinge Mutant (S228P)

[0086] Angal *et al.* (1993) Molecular Immunol. 30, 105-108, have shown that changing Ser 228 in the hinge of a human  $\gamma 4$  chimeric antibody abolishes the formation of half antibody molecules which is a property of natural IgG4 antibodies. Therefore to reduce the heterogeneity of the hENA-2 L235A antibody the S228P change was introduced into the molecule. Mutagenesis was carried by the PCR strand overlap procedure (HO SN *et al.*, 1989) using the following primers:-

R5989

5' CCA TGC CCA TGC CCA GGT AAG CC 3'

R5990

5' CCT GGG CAT GGT GGG CAT GGG GGA CC 3'

[0087] The resulting CDR-grafted heavy chain was combined with the CDR-grafted light chain in a double gene vector for transfection into NSO cells. Cell lines were established producing the hENA-2 L235A, S228P antibody with yields of approximately 400mg/L in suspension culture. The antigen-binding properties of this antibody were indistinguishable from the hENA-2-wt and hENA-2 L235A antibodies. However, the formation of half antibody molecules was reduced from ~10% to less than 1% as judged by SDS PAGE analysis of purified protein.

[0088] CDR-grafted hENA-2 heavy chain vector = pENA216 and double gene expression vector = pENA217 (see Figure 7).

[0089] Figure 16 shows the effects of hENA-2 (L235A) and the above derivative of hENA-2 (L235A, S228P) on the binding of human PMN to E-selectin transfected CHO cells. The method was as described previously on pages 19 to 20. MOPC-21 is used as a negative control antibody.

Referential Examples of therapeutic usesEXAMPLE 2

[0090] We have investigated the effects of ENA-2 F(ab)<sub>2</sub> and the fully engineered human variant, hENA-2 (L235A), in cytokine induced inflammation in baboons.

[0091] Intradermal injections of recombinant human IL-1  $\alpha$  (rhIL-1), 0.1 and 1.0  $\mu$ g/site, and recombinant human TNF  $\alpha$  (rhTNF), 5  $\mu$ g/site, were made in adult baboons. Injections were made, to a pre-determined random grid pattern, at 2, 4 or 8 hours prior to taking full thickness skin biopsies with an 8mm diameter skin biopsy punch. Each mediator was given as a 0.1ml injection in triplicate sites at two time points. Light general anaesthesia was used for the biopsies. The tissues were processed for either formalin fixed paraffin wax histology with haematoxylin and eosin staining or chilled to -70°C for cryostat sectioning and immunohistochemical analysis. Blood samples were taken to determine if the treatment had any effect on circulating blood cells and for antibody pharmacodynamic analysis. Control animals received an equal volume of intravenous saline.

[0092] Injection of rhIL-1 and rhTNF caused an up regulation of E-selectin as detected immunohistochemically and a predominantly neutrophil cellular infiltrate into the deep dermis. Treatment with ENA-2 F(ab)<sub>2</sub> (5mg/kg iv), prior to the cytokine injections attenuated the cellular response at both 4 and 8 hours. This reduction was significant at the 4 hour time point in the TNF injected sites (57% p < 0.05). In a second experiment using the engineered human antibody hENA-2(L235A) (3.0mg/kg iv) there was again a good reduction in cellular inflammation which was significant at 2 and 4 hours for the TNF injected sites (54% and 45% respectively p < 0.05). The degree of inflammation in the IL-1 injected sites was reduced in both the ENA-2 and hENA-2(L235A) treated animals but this did not reach statistical significance. No effect was seen with either antibody on circulating cells.

[0093] The results are shown in Figures 10 to 13.

[0094] This shows that a whole antibody of neutral isotype can be used to inhibit leukocyte extravasation at inflammatory sites. This antibody had a long circulating half-life and could still be detected in the circulation 56 days after treatment (fig. 14) measured as described below.

hENA2(L235A) CELL-BASED ASSAY[0095]AIM

To measure levels of hENA2(L235A) in baboon plasma samples, in order to investigate the Ab's pharmacokinetics.

PRINCIPLE

E-selectin-transfected CHO cells are grown on 96-well tissue culture plates. hENA2(L235A) in baboon plasma binds to E-selectin expressed on cell surfaces and is revealed with a murine anti-human IgG4-HRP & TMB.

REAGENTS

Falcon Microtest III Tissue Culture Plates  
E-selectin-transfected CHO's  
CB2 DMEM Base + 10% Dialysed FCS (tissue culture medium)  
Murine anti-human IgG4-HRP, Serotec MCA517P.  
Dulbecco's PBS  
PBS/1%BSA.  
NMS, Serotec C11 SB.  
TMB substrate.

STANDARD & IAC PREPARATION STANDARD CURVE:

hENA2 diluted to give top standard of 250ng/ml. Doubling dilutions in 1% BSA/PBS to give 125, 62.5, 31.25, 15.6, 7.8 and 3.9 ng/ml (plus zero). Stored in 0.5ml aliquots at -70°C.

IAC's

Controls made up to 60, 25 and 7.5ng/ml hENA2(L235A) in 1% BSA/PBS.

Stored in 0.5ml aliquots at -70°C.

SAMPLE PREPARATION :

Samples diluted in 1% BSA/PBS at a range of dilutions, from 1:10,000 down to 1:10, as for the ELISA.

PROTOCOL:

1. CHO's were plated out into tissue culture plates at 10<sup>5</sup>/ml in tissue culture medium (100  $\mu$ l/well), using only

the middle 60 wells of each plate.

2. Cells were allowed to grow to confluence for 48 hours before assaying, until a monolayer was obtained.

3. Medium was removed with a multichannel pipette and cells were washed once with 100 µl/well PBS (again using a multichannel).

4. Cells were blocked with 1% BSA/PBS, 30 mins, RT, shaking.

5. Plate washed x 2 100 µl/well PBS (with multichannel).

6. 50 µl/well 1% BSA/2% NMS/PBS was added.

7. 50 µl/well standard/IAC/sample, added as appropriate.

8. Plate incubated 1h, RT, shaking gently. Washed x 2 PBS.

9. MaHulG4-HRP diluted at 1:2000, 100 µl/well added.

10. Incubated 30 mins, RT, shaking gently. Washed x 2 PBS as before.

11. 100 µl TMB substrate/well added and absorbance read at 630 nm.

**[0096]** N.B. A multichannel pipette was used to wash the cells and to add each reagent, the plates were shaken gently, they were not tap-dried, and there were only 2 washes with PBS at each stage. These precautions were taken to avoid cells being removed from the plate.

### **EXAMPLE 3**

**[0097]** The engineered human anti-E-selectin antibody, the production of which is described in Example 1 (ie hENA-2,L235A,S228P  $\gamma_4$  isotype) recognises human E-selectin and, under appropriate conditions, prevents PMN adhesion to activated endothelial cells. In order to confirm that the antibody could inhibit the migration of leukocytes into human skin in an *in vivo* setting a model using human skin grafted onto SCID mice has been utilised. This model was first reported by Yan *et al* (J. Clin Invest. 91 986-996 (1993)) and has subsequently been used to show that a murine anti-human E-selectin antibody can inhibit leukocyte trafficking Yan *et al* (J. Immunol 152 3053 (1994)). In this model mouse leukocytes (predominantly PMN) migrate into the human skin via human endothelial cells in response to injected cytokine.

### **METHOD**

**[0098]** After anaesthesia, 6-8 week old SCID mice (purchased from Wistar Institute) were prepared for transplantation by shaving the hair from a 5cm<sup>2</sup> area on each side of the lateral abdominal region. Two circular graft beds approximately 1.5cm in diameter were created on the shaved areas by removing full thickness skin down to the fascia. Full thickness human skin grafts were placed onto the wound beds and held in place with 6-0 non-absorbable monofilament suture material. The human skin consisted of neonatal foreskins from elective circumcisions or normal adult skin removed during plastic surgery. Each experimental mouse received an intravenous injection of 50µg of anti-E-selectin Ab in saline. Control animals received no intravenous injection. The graft on the left side of each animal was then injected with 50µl of normal saline containing a small amount of BSA and colloidal carbon to identify the injection site. The right graft was injected with 6000 units of human recombinant TNF $\alpha$  in 50µl of normal saline containing a small amount of colloidal carbon. After 4 hours the animals were killed and skin sections removed and stored in liquid nitrogen. Immunohistology was performed using standard techniques and sections stained with anti-murine Mac-1, anti-human PECAM-1 and anti-human ICAM-1. This allowed identification of infiltrating murine PMD and confirmed the presence of human endothelial cells lining the blood vessels in the human tissue. The ICAM staining was used as a control to determine that an inflammatory response had been stimulated by the injected TNF. Animals not showing upregulated ICAM-1 expression on basal keratinocytes following TNF injection were excluded from the study. As a further control the saline injected grafts were examined for infiltrating leukocytes. A percentage of animals show high levels of infiltrating leukocytes even in the absence of cytokine injection. This may indicate an ongoing rejection of the graft. Animals showing greater than 100 leukocytes/mm<sup>2</sup> in the saline injected graft were excluded from the study.

**[0099]** Multiple sections (four to six) were cut from the centre of each skin biopsy and three to six randomly chosen x100 microscope fields were examined and the number of infiltrating leukocytes counted.

**[0100]** A total of 14 animals were included in the study but 4 were excluded because the TNF injection site in these

animals showed no ICAM-1 upregulation on basal keratinocytes at the TNF injection site. A further animal was excluded from the study because of high levels of infiltration leukocytes in the saline injected graft. Four animals received 50µg of anti-E-selectin Ab. Four animals received no i.v. treatment.

#### 5 **EXAMPLE 4**

##### **hENA-2(L235A, S228P) — Complement Dependent Lysis**

#### **METHOD**

10 **[0101]** Human dermal microvascular endothelial cells, (HMVECS), (Clonetics) were plated at  $5 \times 10^3$  cells per well into 96 well microtitre plates and cultured in Endothelial Growth medium, (EGM), (Clonetics).

5 Days after plating the cells were labelled with 50Kbq per well  $^{51}$ Chromium overnight at 37°C. The cells were then washed in EGM and treated with human TNF $\alpha$ , (R & D Systems), at 20ng/ml in EGM for 6h at 37°C to upregulate E-Selectin.

15 The plates were washed 5 times in Dulbecco's modified eagle's medium, (DMEM), containing 1% FCS.

Test samples were then added to the appropriate wells. The monoclonal antibodies, (W6/32, mouse anti-human Class 1; hENA-2(L235A) and hENA-2(L235A, S228P) anti-human E-selectin), were added to give a final concentration of 10µg/ml in DMEM + 1%FCS. Rabbit complement (reconstituted according to manufacturer's recommendation) was added to give a final concentration of 12.5% in DMEM + 1% FCS. Where appropriate the complement was heat inactivated at 56°C for 35 min. Maximum lysis was determined by adding 10% sodium dodecyl sulphate, (SDS). After 1h 100µl supernatant was harvested from each well and  $^{51}$ Chromium release in CPM determined on a gamma counter. Results were averages of six replicates. The results are shown in Figure 17.

#### 25 **RESULT**

**[0102]** Neither hENA-2(L235A S228P) nor hENA-2(L235A) mediate complement dependent lysis of human micro-vascular endothelial cells under conditions whereas the mouse monoclonal antibody W6/32 causes lysis.

#### 30 **EXAMPLE 5**

**[0103]** An experiment was carried out to determine the effect of hENA-2(L235A) on endothelial cell integrity. The results are presented in Figure 18 which shows micrographs (x 1890) of haemotoxylin and eosin stained sections from biopsies of baboon skin with cytokine (TNF- $\alpha$ , 5.0µg; 2hr) induced inflammation. Upper micrograph shows endothelium from a saline treated animal with associated PMN infiltrate whilst the lower micrograph shows endothelium from a hENA-2(L235A) (3mg/kg) treated animal with only one or two associated PMNs. No discernible difference in endothelium integrity can be seen between the two treatments.

#### **EXAMPLE 6**

40 **[0104]** Figure 19 shows the inhibition of binding of human neutrophils to human E-selectin transfected CHO cells. The experiment was performed as described in Example 1. The figure shows the effects of different versions of ENA-2 on neutrophil binding from a donor with Fc receptors that interact with many isotypes. The binding is only partly inhibited by the whole murine parent (ENA-2 IG) which is a murine  $\gamma 1$  antibody. The binding can be inhibited by F(ab')<sub>2</sub> of the same antibody showing that most of the binding is caused by an Fc interaction. The chimeric ENA-2 antibody is a full length antibody with the Fc region found in hENA-2. This clearly does not interact with the Fc receptors of this donor and blocks nearly as well as the F(ab')<sub>2</sub> fragment.

#### 50 **Claims**

1. An engineered human antibody having specificity for E-selectin, wherein the complementarity determining regions of the variable domains in said antibody correspond to those of the murine antibody ENA-2 and the following framework residues in said antibody correspond to the same residues of the murine antibody ENA-2:

- heavy chain: 48, 67, 69, 73, 93 & 94
- light chain: 48, 60, 63, 70, 111 & 113

2. An antibody according to claim 1 wherein said antibody is of neutral isotype, **characterised in that** the interactions with antibody Fc receptors and complement are absent or so weak as to cause minimal detrimental physiological effects such as antibody dependent cellular cytotoxicity (ADCC) and/or complement mediated lysis and also the antibody produces a minimal immune response in the host.
3. An antibody according to claim 2, wherein residue 235 in the CH2 domain is alanine.
4. An antibody according to claim 2 or claim 3, being of human  $\gamma 4$  isotype.
5. An antibody according to claim 4, wherein the serine residue at position 228 of the hinge region is a proline residue.
6. An antibody according to any preceding claim, wherein the variable domains in said antibody are those shown in Figures 2a and 2b.
7. A DNA molecule encoding an antibody according to any preceding claim.
8. An antibody according to any one of claims 2-6, linked to a therapeutic agent.
9. An antibody according to any one of claims 2-6, wherein a therapeutic agent is linked to the antibody via a liposome.

#### Patentansprüche

1. Gentechnisch hergestellter, menschlicher Antikörper mit Spezifität für E-Selectin, wobei die hypervariablen Regionen der variablen Domänen in dem Antikörper denen des Maus-Antikörpers ENA-2 entsprechen, und die folgenden Gerüstreste in dem Antikörper denselben Resten des Maus-Antikörpers ENA-2 entsprechen:
  - schwere Kette: 48, 67, 69, 73, 93 & 94
  - leichte Kette: 48, 60, 63, 70, 111 & 113.
2. Antikörper nach Anspruch 1, wobei der Antikörper ein neutraler Isotyp ist, **dadurch gekennzeichnet, daß** die Wechselwirkungen mit Antikörper-Fc-Rezeptoren und des Komplements nicht vorhanden oder so schwach sind, daß minimale schädigende, physiologische Wirkungen wie Antikörper-abhängige zelluläre Cytotoxizität (ADCC) und/oder Komplementvermittelte Lyse verursacht werden, und der Antikörper auch eine minimale Immunantwort im Wirt auslöst.
3. Antikörper nach Anspruch 2, wobei der Rest 235 in der CH2-Domäne Alanin ist.
4. Antikörper nach Anspruch 2 oder Anspruch 3, der ein menschlicher  $\gamma 4$ -Isotyp ist.
5. Antikörper nach Anspruch 4, wobei der Serinrest bei Position 228 der Gelenkregion ein Prolinrest ist.
6. Antikörper nach einem der vorhergehenden Ansprüche, wobei die variablen Domänen in dem Antikörper die in den Figuren 2a und 2b gezeigten sind.
7. DNA-Molekül, das für einen Antikörper nach einem der vorhergehenden Ansprüche kodiert.
8. Antikörper nach einem der Ansprüche 2 bis 6, welcher an einen therapeutischen Wirkstoff gebunden ist.
9. Antikörper nach einem der Ansprüche 2 bis 6, wobei ein therapeutischer Wirkstoff über ein Liposom an den Antikörper gebunden ist.

#### Revendications

1. Anticorps humain produit par génie génétique, ayant une spécificité pour la sélectine E, dans lequel les régions déterminant la complémentarité des domaines variables dudit anticorps correspondent à celles de l'anticorps murin ENA-2 et les résidus suivants de la région charpente dudit anticorps correspondent aux mêmes résidus de l'anti-

corps murin ENA-2:

- chaîne lourde: 48, 67, 69, 73, 93 & 94
- chaîne légère: 48, 60, 63, 70, 111 & 113

5

2. Anticorps selon la revendication 1, dans lequel ledit anticorps est d'isotype neutre, **caractérisé en ce que** les interactions avec les récepteurs Fc d'anticorps et le complément sont absentes ou trop faibles pour provoquer des effets physiologiques nocifs minimum tels qu'une cytotoxicité à médiation cellulaire dépendant des anticorps (ADCC) et/ou une lyse à médiation complémentaire et aussi **en ce que** l'anticorps produit une réponse immunitaire minimale chez l'hôte.

10

3. Anticorps selon la revendication 2, dans lequel le résidu 235 du domaine CH2 est l'alanine.

4. Anticorps selon la revendication 2 ou la revendication 3, qui est d'isotype  $\gamma 4$  humain.

15

5. Anticorps selon la revendication 4, dans lequel le résidu sérine en position 228 de la région charnière est un résidu proline.

6. Anticorps selon l'une quelconque des revendications précédentes, dans lequel les domaines variables dans ledit anticorps sont ceux présentés dans les figures 2a et 2b.

20

7. Molécule d'ADN codant pour un anticorps selon l'une quelconque des revendications précédentes.

8. Anticorps selon l'une quelconque des revendications 2 à 6, lié à un agent thérapeutique.

25

9. Anticorps selon l'une quelconque des revendications 2 à 6, dans lequel un agent thérapeutique est lié à l'anticorps par l'intermédiaire d'un liposome.

30

35

40

45

50

55



**LIGHT CHAIN VARIABLE DOMAIN:**

ATG AGT GTG CCC ACT CAG GTC CTC GGG TTG CTG CTG CTG TGG CTT ACA GAT GCC  
TAC TCA CAC CAC GGC TGA GTC CAG GAC CCC AAC GAC GAC GAC ACC GAA TGT CTA CGG  
M S V P T Q V L G L L L L W L T D A>

AGA TGT GAT ATC GTG ATG ACC CAG TCT CCC TCC TCC CTG ACT GTG ACA GCA GGA  
TCTT ACA CTA TAG CAC TAC TGG GTC AGA GGG AGG AGG GAC TGA CAC TGT CGT CCT  
R C D I V M T Q S P S S L T V T A G>

GAG AAG GTC ACT ATG CGC TGC AAG TCC AGT CAG AGT CTG TTA AAC AGT GGA AAT  
CTC TTC CAG TGA TAC GCG ACG TTC AGG TCA GTC TCA GAC AAT TTG TCA CCT TTA  
E K V T M R C K S S O S L L N S G N>

CAA CAG AAC TAC TTG ACC TGG TAC CAG CAG AAA CCA GGG CAG CCT CCT AAA CTT  
GTT GTC TTG ATG AAC TGG ACC ATG GTC GTC TTT GGT CCC GTC GGA GGA TTT GAA  
Q Q N Y L T W Y Q Q K P G Q P P K L>

TTG ATC TAT TGG GCA TCC ACT AGG GAA TCT GGG GTC CCT GAT CGC TTC ACA GGC  
AAC TAG ATA ACC CGT AGG TGA TCC CTT AGA CCC CAG GGA CTA GCG AAG TGT CCG  
L I Y W A S T R E S G V P D R F T G>

AGT GGA TCT GGA ACA GAT TTC ACT CTC ACC ATC AGC AGT GTG CAG ACT GAA GAC  
TCA CCT AGA CCT TGT CTA AAG TGA GAG TGG TAG TCG TCA CAC GTC TGA CTT CTG  
S G S G T D F T L T I S S V Q T E D>

CTG GCA GTT TAT TAC TGT CAG AAT GAT TAT TAT CCG CTC ACG TTC GGT GCT  
GAC CGT CAA ATA ATG ACA GTC TTA CTA ATA GGC GAG TGC AAG CCA CGA CGA  
I. A V Y Y C Q N D Y D Y P L T F G A>

GGC ACC AAG CTG GAG ATC AAA GGT  
CCG TGG TTC GAC CTC TAG TTT GCA  
G T K L E I K R>

FIG. 1a

## HEAVY CHAIN VARIABLE DOMAIN.

CCG CCA CC ATG GGA TGG AGC TGG ACC TAA TTT ATC TTC CTC CTC TCA GTA ATT  
 GGC GGT GG TAC CCT ACC TCG ACC TAA AAA TAG AAG GAG GAG AGT CAT TAA  
 M G W S W I F I F L L S V I>  
 E G V  
 G G V

GCA GGT GTC CAA TCC CAG GTT CAA CTG CAG CAG TCT GGG ACT GAA CTG GTG  
 CGT CCA CAG GTT AGG GTC CAA GTT GAC GTC AGA CCC TGA CTT GAC CAC  
 A G V Q S Q V Q L Q Q S G T E L V>

AGG CCT GGG GCT TCA GTG ACG CTG TCC TGC AAG GCT TCG GGC TAC ACA TTT  
 TCC GGA CCC CGA AGT CAC TGC GAC AGG ACG TTC CGA AGC CCG ATG TGT AAA  
 R P G A S V T L S C K A S G Y T F

ACT GAC CAT GAA ATG CAC TGG GTG AAG CAG ACA CCT GTG CTT GGC CTG GAA  
 TGA CTG GTA CTT TAC GTG ACC CAC TTC GTC TGT GGA CAC GAA CCG GAC CTT  
 T D H E M H W V K Q T P V L G L E>

TGG ATT GGA ACT ATT GAT CCT GAA ACT GGT GGT ACT GCC TAC AAT CAG AAG  
 ACC TAA CCT TGA TAA CTA GGA CTT TGA CCA CCA TGA CGG ATG TTA GTC TTC  
 W I G T I D P E T G G T A Y N Q K

TTC AAG GGC AAG GCC ACA CTG ACT GCA GAC AAA TCC TCC ACT ACA GCC TAC  
 AAG TTC CCG TTC CGG TGT GAC TGA CGT CTG TTT AGG AGG TGA TGT CGG ATG  
 F K G K A T L T A D K S S T T A Y>

ATG GAC CTC CGC GGC CTG ACA TCT GAG GAC TCT GCC GTC TTT TAC TGT ACA  
 TAC CTG GAG GCG CCG GAC TGT AGA CTC CTG AGA CGG CAG AAA ATG ACA TGT  
 M D L R G L T S E D S A V F Y C T

GTC CTA AGG ATG GAC TAC TGG GGT CAA GGA ACC TCA CTC ACA GTC TCC GCA  
 CAG GAT TCC TAC CTG ATG ACC CCA GTT CCT TGG AGT GAG TGT CAG AGG CGT  
 V L R M D Y W G Q G T S L T V S A

FIG.1b

## FIG. 2a

## Heavy chain variable domain.

```

GCG CGC AAG CTT GCC GCC ACC ATG GAA TGG AGC TGG GTC TTT CTC TTC
CGC GCG TTC GAA CGG CGG TGG TAC CTT ACC TCG ACC CAG AAA GAG AAG
      M   E   W   S   W   V   F   L   F>

TTC CTG TCA GTA ACT ACA GGA GTC CAT TCT CAG GTG CAG CTG GTG CAG
AAG GAC AGT CAT TGA TGT CCT CAG GTA AGA GTC CAC GTC GAC CAC GTC
  F   L   S   V   T   T   G   V   H   S   Q   V   Q   L   V   Q>

TCT GGA GCA GAG GTG AAG AAG CCT GGA TCT TCT GTG AAG GTG TCT TGT
AGA CCT CGT CTC CAC TTC TTC GGA CCT AGA AGA CAC TTC CAC AGA ACA
  S   G   A   E   V   K   K   P   G   S   S   V   K   V   S   C>

AAG GCA TCT GGA TAC ACA TTC ACA GAC CAC GAG ATG CAC TGG GTG AGA
TTC CGT AGA CCT ATG TGT AAG TGT CTG GTG CTC TAC GTG ACC CAC TCT
  K   A   S   G   Y   T   F   T   D   H   E   M   H   W   V   R>

CAG GCA CCT GGA CAG GGA CTC GAG TGG ATT GGA ACA ATT GAC CCT GAG
GTC CGT GGA CCT GTC CCT GAG CTC ACC TAA CCT TGT TAA CTG GGA CTC
  Q   A   P   G   Q   G   L   E   W   I   G   T   I   D   P   E>

ACA GGA GGA ACA GCC TAC AAT CAG AAG TTC AAG GGA AGA GCA ACA CTG
TGT CCT CCT TGT CGG ATG TTA GTC TTC AAG TTC CCT TTT CGT TGT GAC
  T   G   G   T   A   Y   N   Q   K   F   K   G   R   A   T   L>

ACA GCA GAC AAG TCT ACG AAT ACC GCC TAC ATG GAG CTG TCT TCT CTG
TGT CGT CTG TTC AGA TGC TTA TGG CGG ATG TAC CTC GAC AGA CCT GAC
  T   A   D   K   S   T   N   T   A   Y   M   E   L   S   S   L>

AGA TCT GAG GAC ACA GCA GTG TAC TAC TGT ACA GTG CTC AGA ATG GAC
TCT AGA CTC CTG TGT CGT CAC AAG ATG ACA TGT CAC GAG TCT TAC CTG
  R   S   E   D   T   A   V   Y   Y   C   T   V   L   R   M   D>

TAC TGG GGA CAG GGA ACA CTG GTG ACA GTG TCT TCT
ATG ACC CCT GTC CCT TGT GAC CAC TGT CAC AGA AGA
  Y   W   G   Q   G   T   L   V   T   V   S   S

```

## FIG. 2b

Light chain variable domain.

```

GGA CTG TTC GAA GCC GCC ACC ATG TCT GTC CCC ACC CAA GTC CTC GGA
CCT GAC AAG CTT CGG CGG TGG TAC AGA CAG GGG TGG GTT CAG GAG CCT
      M   S   V   P   T   Q   V   L   G>

CTC CTG CTG CTG TGG CTT ACA GAT GCC AGA TGC GAT ATC CAG ATG ACT
GAG GAC GAC GAC ACC GAA TGT CTA CGG TCT ACG CTA TAG GTC TAC TGA
  L   L   L   L   W   L   T   D   A   R   C   D   I   Q   M   T>

CAG AGT CCA AGT ACT CTC AGT GCC AGT GTA GGT GAT AGG GTC ACC ATC
GTC TCA GGT TCA TGA GAG TCA CGG TCA CAT CCA CTA TCC CAG TGG TAG
  Q   S   P   S   T   L   S   A   S   V   G   D   R   V   T   I>

ACT TGT AAG TCT TCT CAA TCT CTC TTA AAC TCC GGT AAC CAG CAG AAC
TCC ACA TTC AGA AGA GTT AGA GAG AAT TTG AGG CCA TTG GTC GTC TTG
  T   C   K   S   S   Q   S   L   L   N   S   G   N   Q   Q   N>

TAC CTC ACT TGG TAC CAG CAG AAA CCA GGT AAA GCC CCA AAG CTC CTC
ATG GAG TGA ACC ATG GTC GTC TTT GGT CCA TTT CGG GGT TTC GAG GAG
  Y   L   T   W   Y   Q   Q   K   P   G   K   A   P   K   L   L>

ATC TAT TGG GCC TCT ACT AGG GAA TCT GGT GTA CCA GAT AGA TTC ACT
TAG ATA ACC CGG AGA TGA TCC CTT AGA CCA CAT GGT CTA TCT AAG TGA
  I   Y   W   A   S   T   R   E   S   G   V   P   D   R   F   T>


GGT AGT GGT AGT GGT ACT GAT TTC ACT CTC ACT ATC AGT AGT CTC CAG
CCA TCA CCA TCA CCA TGA CTA AAG TGA GAG TGA TAG TCA TCA GAG GTC
  G   S   G   S   G   T   D   F   T   L   T   I   S   S   L   Q>

CCA GAT GAT TTC GCC ACT TAT TAC TGT CAG AAC GAT TAC GAT TAC CCA
GGT CTA CTA AAG CGG TGA ATA ATG ACA GTC TTG CTA ATG CTA ATG GGT
  P   D   D   F   A   T   Y   Y   C   Q   N   D   Y   D   Y   P>

TTA ACT TTC GGT CAG GGT ACT AAA GTA GAA ATC AAA CGT
AAT TGA AAG CCA GTC CCA TGA TTT CAT CTT TAG TTT CCA
  L   T   F   G   Q   G   T   K   V   E   I   K   R

```

FIG. 3a






FIG. 3b

FIG. 4a





FIG. 4b

## ADHESION ASSAY





G1	A P E L L G G P S V F L F P P K D T.....
G2	A P P V A G . P S V F L F P P K D T.....
G3	A P E L L G G P S V F L F P P K D T.....
G4	A P E F L G G P S V F L F P P K D T.....

[b] Sequences of mutagenic oligonucleotides.

[illegible]

## Oligonucleotides

AC

5' CCT GAG TTC GTC GGG GGA CCA TCA GTC TTC 3'

GA AGG AGT CGT GGA CTC AAG CAG CCC CCT GG

TG

FIG. 6




FIG. 7






FIG. 8





Polylinker 1: 2.56 / NotI, BamHI, NruI, NaeI

FIG. 9





FIG. 10






FIG. 11

FIG. 12






FIG. 13





FIG. 14






FIG. 15

\* DIFFERENT FROM THE OTHER THREE GROUPS BY ANOVA ( $P < 0.05$ )




FIG. 16




FIG. 17




FIG. 18




A



B

FIG. 19

## ADHESION ASSAY



**THIS PAGE BLANK (USPTO)**